

TRULICITY

BY SUKAINA DAMJI

Trulicity is indicated for the treatment of adults with insufficiently controlled type 2 diabetes mellitus as an adjunct to diet and exercise

Monotherapy: The recommended dose is 0.75 mg once weekly.
Add-on therapy: The recommended dose is 1.5 mg once weekly.
For additional glycaemic control: • the 1.5 mg dose may be increased after at least 4 weeks to 3 mg once weekly. • the 3 mg dose may be increased after at least 4 weeks to 4.5 mg once weekly.
The maximum dose is 4.5 mg once weekly. .

Trulicity is to be injected subcutaneously in the abdomen, thigh or upper arm.

The dose can be administered at any time of day, with or without meals.

The day of weekly administration can be changed if necessary, as long as the last dose was administered 3 or more days (72 hours) before.

Possible side effects; hypoglycemia, decreased appetite , nausea, diarrhoea, flatulence, acidity.


Store in a refrigerator (2 °C – 8 °C). In-use Trulicity may be stored unrefrigerated for up to 14 days at a temperature not above 30 °C.

Trulicity 1.5 mg solution for injection in pre-filled pen
dulaglutide




- The pen is a disposable, pre-filled delivery device that is ready to use. Each pen contains one weekly dose of Trulicity (1.5 mg). Each pen delivers one dose only.
- • Trulicity is administered once a week. You may want to mark your calendar to remind you when to inject your next dose.
- • When you press the green injection button, the pen will automatically insert the needle into your skin, inject the medicine, and pull back (retract) the needle after the injection is complete.

1. **UNCAP**
2. **PLACE AND UNLOCK**
3. **PRESS AND HOLD**




1 UNCAP

-  Make sure the pen is **locked**.
- Pull off the grey base cap.

Do not put the base cap back on – this could damage the needle. Do not touch the needle.

- When removed, the grey base cap appears as shown and can be discarded.




If the grey base cap does not look like the image above:

- **Do not** use the Pen.
- Safely store the Pen and cap and contact **Lilly**.
- Use a new Pen.

2 PLACE AND UNLOCK


- Place the clear base flat and firmly against your skin at the injection site.

-  **Unlock** by turning the lock ring.




3 PRESS AND HOLD

- Press and hold the green injection button; you will hear a loud click.

-  Continue holding the clear base firmly against your skin until you hear a second click. This occurs when the needle starts retracting in about 5-10 seconds.

- Remove the pen from your skin.



- You will know your injection is complete when the grey part is visible.